

# **Spatial and temporal dynamics of photosynthesis regulation of genetically defined coral/algal symbiosis associations**

A thesis submitted in fulfilment of the requirements for the  
degree of Doctor of Philosophy by

**Karin Elizabeth Ulstrup**

(B.Sc., M.Sc. Biology, University of Copenhagen)

Institute of Water and Environmental Resource Management  
Department of Environmental Sciences  
University of Technology, Sydney

June 2007

## Certificate

The work presented in this thesis, and the research to which it pertains, are the product of my own work and, to the best of my knowledge and belief, original. Any ideas or quotations from the work of others, published or otherwise, are fully acknowledged in accordance with the standard referencing practices of the discipline. Co-authors of published, submitted articles and articles in preparation were involved in two or more stages of a study including **a)** planning of the study, **b)** essential expertise and support with field and laboratory work, **c)** critical evaluation of the material and literature and **d)** final interpretation and writing up of the results. Material has not been submitted, either in whole or in part, for a degree at this or any other university.




Signed \_\_\_\_\_

Karin E. Ulstrup (PhD Candidate)



Institute for Water  
& Environmental  
Resource Management



Australian Institute of Marine Science

Headquarters - Cape Ferguson, Queensland



## Acknowledgements

Facilities for my PhD studies were provided by University of Technology, Sydney (UTS), Australian Institute of Marine Science (AIMS), University of Sydney (USYD) and the University of Copenhagen, Denmark (KU). These places all provided expertise and infrastructure for my project.

I would like to thank my principal supervisor, Assoc. Prof. Peter Ralph (UTS) for his support and know-how throughout the duration of my candidature. Dr. Madeleine van Oppen (AIMS) provided invaluable guidance on the molecular part of my project and personal encouragement. I wish to thank Prof. Michael Kühl (KU) for inspiration and for teaching me the principles of working with microsensors. Prof. Tony Larkum (USYD) and Dr. Ray Berkelmans (AIMS) were also instrumental in the success of parts of my thesis for which I am sincerely grateful. All of the above provided stimulating discussions and assisted in clarifying and refining my thoughts on physiological and genetic aspects of coral science.

The field work in this study was undertaken at Heron Island Research Station, Lizard Island Research Station and on vessels belonging to the AIMS fleet. My time spent in the field was both productive and enjoyable as a result of the thoughtfulness, advice and efforts of the staff at these facilities. I would like to thank the following for their hands-on assistance in the field: Jeremy Downs, Steve Hornabrook, Martin Macnaughton, Tim Cooper and Jens Ulstrup who all volunteered their time and assistance. In the laboratory I was assisted by Martine Goldner, Alex Griffin, Lesa Peplow and Jeremy Downs. I also thank all members of the Aquatic Photosynthesis Lab (UTS). Special thanks to my family who provided me with personal support and encouragement.

Funding from a number of organisations made this project possible and I am grateful for their contribution. University of Technology, Sydney awarded me an International Postgraduate Research Scholarship and a Science Faculty stipend. Research support was generously provided by the National Geographic Society, PADI Foundation, Great Barrier Reef Marine Park Authority, HRH Crown Prince Frederiks Foundation, Denmark, Winifred Violet Scott Foundation, Sigma Xi and a Graduate Fellowship from the International Society for Reef Studies. The Australian Coral Reef Society, Knud Højgaards Fond, Denmark, as well as internal research allowances at

University of Technology, Sydney provided me with financial support to attend scientific conferences.



## **Publications**

### Peer-reviewed journal articles arising directly from this thesis

**Ulstrup KE**, Berkelmans R, Ralph PJ, van Oppen MJH (2006) Variation in bleaching sensitivity of two coral species across a latitudinal gradient on the Great Barrier Reef: the role of zooxanthellae. *Marine Ecology Progress Series* 314:135-148.

**Ulstrup KE**, Ralph PJ, Larkum AWD, Kühl M (2006) Intra-colonial variability in light acclimation of zooxanthellae in coral tissues of *Pocillopora damicornis*. *Marine Biology* 149:1325-1335.

### Peer-reviewed journal articles relevant to the thesis but not forming part of it

**Ulstrup KE**, Kühl M, Bourne D (2007) Zooxanthellae harvested by ciliates associated with brown band syndrome of corals remain photosynthetically competent *Applied Environmental Microbiology* 73:1968-1975

**Ulstrup KE**, Hill R, Ralph PJ (2005) Photosynthetic impact of hypoxia on *in hospite* zooxanthellae in the scleractinian coral *Pocillopora damicornis*. *Marine Ecology Progress Series* 286:125-132.

### Conference abstracts (\* poster)

**Ulstrup KE**, Ralph PJ, Larkum AWD, Kühl M (2006) Intra-colonial variability in light acclimation of zooxanthellae in coral tissues. European Meeting for International Society for Reef Studies, Bremen.

**\*Ulstrup KE**, Ralph PJ, Larkum AWD, Kühl M (2006) Intra-colonial variability in light acclimation of zooxanthellae in coral tissues. American Society for Limnology and Oceanography, 6<sup>th</sup> Ocean Sciences Meeting, Honolulu, USA.

**Ulstrup KE**, Ralph PJ, Larkum AWD, Kühl M (2005) Intra-colonial variability in light acclimation of zooxanthellae in coral tissues. Australian Coral Reef Society 81<sup>st</sup> Annual Conference, Heron Island, Australia.

Hill R, **Ulstrup KE** (2005) Imaging-PAM: operation and possibilities. Coral Reef Targeted Research and Capacity Building for Management Workshop, Puerto Morelos, Mexico.

**Ulstrup KE** (2004) Coral Bleaching - Integrating physiology with genetics. Scandinavian Protozoological Meeting, Helsingør, Denmark.

**Ulstrup KE**, Berkelmans R, Ralph P, van Oppen M (2004) Bleaching sensitivity of two conspecific corals on a latitudinal gradient. International Society for Reef Studies, 10<sup>th</sup> International Coral Reef Symposium, Okinawa, Japan.

## Table of contents

Certificate .....	I
Acknowledgements .....	II
Publications .....	IV
Table of contents .....	VI
Table of figures .....	IX
Table of tables .....	XIII
Abstract .....	XVIII

## Chapter 1 Background, concepts and thesis outline ..... 1

1.0 Background .....	1
1.1 Thermal sensitivity of corals and global warming .....	1
1.2 The role of zooxanthellae .....	3
1.2.1 Diversity, occurrence and distribution of <i>Symbiodinium</i> .....	3
1.2.2 Physiological diversity of zooxanthellae .....	4
1.2.3 Photosynthetic controls of zooxanthellae <i>in hospite</i> .....	6
1.3 Concept of scales .....	8
1.4 Thesis outline .....	8

## Chapter 2 Intra-colonial variability in light acclimation of zooxanthellae in coral tissues of *Pocillopora damicornis* ..... 10

2.0 Introduction .....	10
2.1 Materials and methods .....	13
2.1.1 Sampling and experimental setup .....	13
2.1.2 Scalar irradiance .....	14
2.1.3 Measuring sequence .....	14
2.1.4 Curve fitting .....	15
2.1.5 Statistical analysis .....	16
2.2 Results .....	16
2.3 Discussion .....	23
2.3.1 Light microclimate .....	24
2.3.2 Oxygen dynamics .....	25
2.3.3 Steady-state relationships of gross photosynthesis rate, O <sub>2</sub> concentration and rETR .....	26
2.4 Conclusion .....	28

## Chapter 3 Inter-polyp genetic and physiological characterisation of *Symbiodinium* in an *Acropora valida* colony ..... 29

3.0 Introduction .....	29
3.1 Methods and materials .....	31
3.1.1 DNA extraction and assessment of zooxanthella diversity and dominance .....	32
3.1.2 Experimental setup .....	32
3.1.3 Measuring sequence .....	33
3.1.4 Curve fitting .....	34
3.1.5 Statistical analysis .....	34
3.2 Results .....	35

3.2.1 Genetic composition and distribution of <i>Symbiodinium</i> .....	35
3.2.2 Photo-physiology of individual polyps with contrasting branch orientation and symbiont composition .....	36
3.3 Discussion .....	38
3.3.1 Relative occurrence of clade A .....	39
3.3.2 Photo-physiology of individual polyps with contrasting orientation and symbiont composition .....	40
3.3.2.1 Oxygen dynamics.....	40
3.3.2.3 Influence of light history on capacity for photo-acclimation.....	41

## **Chapter 4 Seasonal variation in photo-physiological functions in *Symbiodinium* communities in two scleractinian corals .....**

43

4.0 Introduction .....	43
4.1 Materials and methods .....	45
4.1.1 Annual temperature.....	45
4.1.2 Molecular methods.....	45
4.1.3 Symbiont cell determination .....	46
4.1.4 Zooxanthellar xanthophyll determination.....	46
4.1.5 PAM fluorescence measurements of PSII.....	47
4.1.6 Fast induction kinetics of PSII.....	48
4.1.7 Statistical analyses .....	49
4.2 Results .....	49
4.2.1 Measurements of temperature .....	49
4.2.2 Genotype variability .....	50
4.2.3 Symbiont cell density .....	51
4.2.4 Xanthophyll cycling and NPQ <sub>RLC</sub> .....	52
4.2.5 Quantum yields .....	53
4.2.6 Descriptive parameters of fitted rETR-curves .....	55
4.2.7 Fast induction kinetics.....	56
4.3 Discussion .....	57
4.3.1 Genetic diversity of <i>Symbiodinium</i> .....	58
4.3.2 Symbiont cell density .....	59
4.3.3 Photo-protection.....	59
4.3.4 Photochemical efficiency and photo-acclimation .....	60
4.4 Conclusions .....	61

## **Chapter 5 Variation in photosynthetic performance of conspecific corals from two geographically distinct populations .....**

63

5.0 Introduction .....	63
5.1.1 Experimental procedure .....	67
5.1.2 Curve fitting .....	69
5.1.3 Genetic characterisation of zooxanthellae .....	69
5.1.4 Statistical analysis .....	70
5.2 Results .....	70
5.2.1 Environmental measurements .....	70
5.2.2 Physiological measurements .....	71
5.2.2.1 Comparisons between sampling times coinciding with winter and summer .....	71
5.2.2.2 Effects of temperature .....	72



5.2.2.3 Effects of location .....	72
5.2.2.4 Interactions among temperatures and between locations.....	72
5.2.3 Genetic characterisation of zooxanthellae .....	77
5.3 Discussion .....	77
5.3.1 Effects of sampling time .....	77
5.3.2 Temperature and location.....	78
5.3.3 Summary .....	79

## **Chapter 6 Variation in bleaching sensitivity of two coral species across a latitudinal gradient on the Great Barrier Reef: the role of zooxanthellae ..... 80**

6.0 Introduction.....	80
6.1 Materials and methods .....	82
6.1.1 Sampling .....	82
6.1.2 Sea temperature measurements .....	84
6.1.3 Experimental procedure .....	85
6.1.4 Genetic identification of zooxanthellae in field colonies.....	87
6.1.5 Statistical analyses .....	88
6.2 Results.....	88
6.2.1 Temperature range .....	88
6.2.2 Mortality.....	89
6.2.3 Zooxanthella density .....	91
6.2.4 Photochemical efficiency ( $F_v/F_m$ ) .....	94
6.2.4.1 <i>Pocillopora damicornis</i> .....	94
6.2.4.2 <i>T. reniformis</i> .....	96
6.2.5 Genotyping of zooxanthellae .....	98
6.3 Discussion .....	98
6.3.1 Local adaptation and phenotypic plasticity.....	99
6.3.2 Mortality.....	100
6.3.3 Zooxanthella density .....	101
6.3.4 Photochemical efficiency .....	101
6.3.5 Symbiont selection .....	102
6.4 Conclusions.....	103

## **Chapter 7 Key findings, implications and future research ..... 104**

7.0 General discussion .....	104
7.0.1 Technical points on photo-physiological measurements at the microscale .....	105
7.0.2 Specificity of coral/algal associations.....	106
7.0.3 Mixed coral/algal associations .....	107
7.0.4 Large-scale studies of coral photobiology in time and space .....	108
7.1 Summary of key findings .....	109
7.2 Future research .....	110

References.....	112
Appendix A .....	129
Appendix B .....	131
Appendix C .....	134
Appendix D.....	149
Appendix E .....	156

## Table of figures

Figure 2.1. Spectral scalar irradiance measurements of sun- and shade-adapted polyp tissues of *P. damicornis*. Data are normalised to the incident downwelling scalar irradiance ( $E_d$ ). Arrow indicates the absorption wavelength of chlorophyll *a* (675 nm).

..... 17

Figure 2.2a-b. Steady-state levels of  $O_2$  concentration at the surface of sun- (○) and shade-adapted (●) coenosarc a) and polyp b) tissues of *P. damicornis* ( $n = 5 \pm SE$ ) in response to increasing scalar irradiance. Arrows indicate the compensation irradiance ( $E_c$ ), above which the oxygen level at the tissue surface exceeds the level of the overlaying air saturated seawater ( $208 \mu\text{mol l}^{-1}$ )..... 18

Figure 2.3a-d. Steady-state rETR (●) and gross photosynthesis rate (bars) as a function of increasing scalar irradiance ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) of a) sun- and b) shade-adapted coenosarc and (c) sun- and (d) shade-adapted polyp tissues of *P. damicornis* ( $n = 5 \pm SE$ ). The fitted rETR curve is superimposed with a solid line. The fitted gross photosynthesis rate curve is superimposed with a broken line. .... 20

Figure 2.4a-b. Steady-state rETR and gross photosynthesis rate of sun- (○) and shade-adapted (●) of a) coenosarc and b) polyp tissues of *P. damicornis* in response to increasing scalar irradiance ( $n = 5 \pm SE$ ). .... 21

Figure 2.5a-b. Gross photosynthesis rate and  $O_2$  concentration of sun- (○) and shade-adapted (●) a) coenosarc and b) polyp tissues of *P. damicornis* in response to increasing scalar irradiance ( $n = 5 \pm SE$ ).  $r^2$ , correlation coefficient; a, slope of the correlation line. .... 22

Figure 2.6. Steady-state rETR of sun- (○) and shade-adapted (●) of a) coenosarc and b) polyp tissues of *P. damicornis* as a function of  $O_2$  concentration ( $n = 5 \pm SE$ )..... 23

Figure 3.1a-b. Steady-state light curves of  $O_2$  concentration ( $\mu\text{mol l}^{-1} \pm SE$ ) at the surface of a) sun- (○) and shade-adapted (●) polyps, b) clade A + C (○) and clade C polyps (●) of *A. valida*. The horizontal line indicates the level of the air-saturation at the surface of the coral ( $230.9 \mu\text{mol l}^{-1}$ ) and the vertical arrows indicate the compensation irradiance,  $E_c$ , estimate. Significant difference ( $p = 0.002$ ) is denoted with asterisks (\*). .... 36

Figure 3.2a-d. Steady-state relative electron transport rate (rETR) (●) and gross photosynthesis rate (white bars) as a function of increasing irradiance ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) of individual polyps of *A. valida* for a) sun-adapted polyps, b) shade-adapted polyps, c) clade A + C polyps and d) clade C polyps. The fitted rETR curves are superimposed with a solid line. The fitted gross photosynthesis rate curves are superimposed with a broken line..... 38

Figure 4.1a-b. Maximum daily temperature (●) at Heron Island and symbiont cell density ( $10^6 \text{ cm}^{-2}$ ) of a) *P. damicornis* and b) *A. valida* in sun- (white bars) and shade-adapted surfaces (hatched bars) of colonies. Averages  $\pm SE$  are shown ( $n = 10$ )..... 52

Figure 4.2a-b. Xanthophyll ratio (ratio of diatoxanthin to the sum of diatoxanthin and diadinoxanthin) in sun- (white bars) and shade-adapted surfaces (hatched bars) and non-



photochemical quenching in sun- (○) and shade-adapted surfaces (●) of individual colonies of a) *P. damicornis* and b) *A. valida*. Averages  $\pm$  SE are shown (n =10).....53

Figure 4.3a-b. Effective quantum yield,  $\Phi_{PSII}$ , measured under quasi-darkness conditions in sun- (○) and shade-adapted (●) surfaces at dusk in a) *P. damicornis* and b) *A. valida*. Averages  $\pm$  SE are shown (n =10).....54

Figure 4.4a-d. Maximum quantum yield ( $F_v/F_m$ ) of zooxanthellae in sun- (○) and shade-adapted (●) surfaces of *P. damicornis* at a) dusk and b) dawn and in *A. valida* at c) dusk and d) dawn. Averages  $\pm$  SE are shown (n =10).....55

Figure 4.5a-h. Amplitude of the O, J, I and P steps (●, ■, ▲ and ▼, respectively) along FICs in *P. damicornis*; a) sun- and b) shade-adapted surfaces and *A. valida*; c) sun- and d) shade-adapted surfaces between January 2004 and July 2006 (n =10  $\pm$  SE). Average FICs (n =10) from *P. damicornis* on e) sun- and f) shade-adapted surfaces and of *A. valida* on g) sun- and h) shade-adapted surfaces are also shown as examples of how the O, J, I and P step amplitudes were obtained. The dashed arrows correlate to the amplitude of each of the steps for these particular FICs.....57

Figure 5.1. Map of the Australian north eastern coast showing collection sites of *Pocillopora damicornis* and *Turbinaria reniformis* and their associated symbiont type. ....65

Figure 5.2. Average daily temperatures for the reef slope (6-9 m depth) for each day (48 measurements per day) at Lizard Island and Davies Reef between May 2005 and April 2006.....66

Figure 5.3. Annual variation in day length (h) at Lizard Island (●) and Broadhurst Reef (○), also representing Davies Reef between May 2005 and April 2006. The data is derived from the Geoscience Australia and represent monthly averages of daily recording  $\pm$  SE. Arrows indicate twice yearly sampling time at Lizard Island (upward-pointing arrows) and Broadhurst Reef and Davies Reef (downward-pointing arrows)..66

Figure 5.4. Descriptive parameters ( $P_{max}$  [ $\text{nmol cm}^{-3} \text{s}^{-1}$ ],  $\alpha$ , and  $E_k$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ]) of fitted PI-curves of *P. damicornis* from Lizard Island (black bars) and Broadhurst Reef in winter and Davies Reef in summer in grey bars as a function of temperature treatment (n =4  $\pm$  SE). Significant differences (p <0.05) are denoted with asterisks (\*) for 2-way ANOVA and with + for 1-way ANOVA. ....73

Figure 5.5. Descriptive parameters ( $P_{max}$  [ $\text{nmol cm}^{-3} \text{s}^{-1}$ ],  $\alpha$ , and  $E_k$  [ $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ]) of fitted PI-curves of *T. reniformis* from Lizard Island (black bars) and Davies Reef (grey bars) as a function of temperature treatment (n =4  $\pm$  SE). Significant differences (p <0.05) are denoted with asterisks (\*) for 2-way ANOVA and with + for 1-way ANOVA. ....74

Figure 5.6. Measurements of  $P_{gmax}$  ( $\text{nmol cm}^{-3} \text{s}^{-1}$ ),  $P_{nmax}$  ( $\text{nmol cm}^{-2} \text{s}^{-1}$ ),  $R_D$  ( $\text{nmol cm}^{-2} \text{s}^{-1}$ ),  $P_{nmax}:R$  and  $O_{2max}$  ( $\mu\text{mol l}^{-1}$ ) for *P. damicornis* in winter at Lizard Island (black bars) and Broadhurst Reef (grey bars) and in summer at Lizard Island (black bars) and Davies Reef (grey bars) (n =4  $\pm$  SE). All maximum measurements are obtained at 900

$\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Significant differences ( $p < 0.05$ ) are denoted with asterisks (\*) for 2-way ANOVA. .... 75

Figure 5.7. Measurements of  $P_{g\text{max}}$  ( $\text{nmol cm}^{-3} \text{ s}^{-1}$ ),  $P_{n\text{max}}$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $R_D$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $P_{n\text{max}}:R$  and  $O_{2\text{max}}$  ( $\mu\text{mol l}^{-1}$ ) for *T. reniformis* in winter and summer at Lizard Island (black bars) and Davies Reef (grey bars) ( $n = 4 \pm \text{SE}$ ). All maximum measurements are obtained at  $900 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Significant differences ( $p < 0.05$ ) are denoted with asterisks (\*) for 2-way ANOVA and with + for 1-way ANOVA. .... 76

Figure 6.1. Map of the Australian eastern sea border showing collection sites: <sup>a</sup>Locations of sea temperature measurements; <sup>b</sup>locations of experimental bleaching samples. .... 83

Figure 6.2. Average daily temperatures for the reef slope (6-9 m depth) for each day (48 measurements per day) at Lizard Island, Davis Reef, Hardy Reef, East Cay and Heron Island. A 10 day smoothing function is applied to indicate the general trend in yearly temperatures. .... 85

Figure 6.3. Accumulated mortality (% of  $n = 27$ ) of explants of *P. damicornis* from Lizard Island (black bars), Big Broadhurst Reef (light grey bars) and Wistari Reef (dark grey bars) over the course of the experiment. Elevated temperature exposure is represented in the first 2 weeks (1-13 days, grey top bar) and recovery ( $26^\circ\text{C}$ ) (15-27 days, white top bar) is represented in the last two weeks. .... 90

Figure 6.4. Accumulated mortality (% of  $n = 27$ ) of explants of *T. reniformis* from Lizard Island (black bars), Big Broadhurst Reef (light grey bars) and Heralds Prong Reef (dark grey bars) over the course of the experiment. Exposure to  $33^\circ\text{C}$  is represented in the first 2 weeks (1-13 days, grey top bar) and recovery ( $26^\circ\text{C}$ ) is represented in the last two weeks (15-27 days, white top bar). .... 91

Figure 6.5. Relative change (%) of zooxanthella density ( $10^6 \text{ cm}^{-2}$ ) in relation to controls of explants of *P. damicornis* from Lizard Island (black bars), Big Broadhurst Reef (light grey bars) and Wistari Reef (dark grey bars) after 2 weeks of exposure to elevated temperatures ( $29^\circ\text{C}$  and  $31^\circ\text{C}$ ) and 2 weeks of recovery. Significant differences were calculated by testing between sites at specific temperatures. Averages are shown ( $n = 3$ ) including SE bars. Significant differences ( $p < 0.05$ ) are shown with an asterisk. .... 93

Figure 6.6. Relative change (%) of zooxanthella density ( $10^6 \text{ cm}^{-2}$ ) in relation to controls of explants of *T. reniformis* from Lizard Island (black bars), Big Broadhurst Reef (light grey bars) and Heralds Prong Reef (dark grey bars) after 2 weeks of exposure to elevated temperatures ( $29^\circ\text{C}$  and  $31^\circ\text{C}$ ) and 2 weeks of recovery ( $n = 3 \pm \text{SE}$ ). Significant differences were calculated by testing between sites at specific temperatures. Significant differences ( $p < 0.05$ ) are denoted with asterisk. .... 94

Figure 6.7a-d. Relative change (%) of photochemical efficiency ( $F_v/F_m$ ) in relation to pre-bleaching responses ( $n = 27 \pm \text{SE}$ ) of explants of *P. damicornis* from Lizard Island (●), Big Broadhurst Reef (○) and Wistari Reef (▼). Two weeks of control (a: $26^\circ\text{C}$ ) and bleaching treatments (b: $29^\circ\text{C}$ , c: $31^\circ\text{C}$ , d: $33^\circ\text{C}$ ) are shown (1-13 days, grey top bars) followed by 2 weeks (15-27 days, white top bars) of recovery at  $26^\circ\text{C}$ . .... 96



Figure 6.8a-d. Relative change (%) of photochemical efficiency ( $F_v/F_m$ ) in relation to pre-bleaching responses ( $n = 27 \pm SE$ ) of explants of *T. reniformis* from Lizard Island (●), Big Broadhurst Reef (○) and Heralds Prong Reef (▼). Two weeks of control (a:26°C) and bleaching treatments (b:29°C, c:31°C, d:33°C) are shown (1-13 days, grey top bars) followed by 2 weeks (15-27 days, white top bars) of recovery at 26°C.....97

Appendix A, Figure 1. Scheme of experimental setup for combined microsensor measurements of oxygen and variable chlorophyll *a* fluorescence. The combined microsensor is mounted on a micromanipulator (MM). The O<sub>2</sub> microelectrode part of the sensor is connected to a picoamperemeter (pA) and the sensor signal is recorded on a strip chart recorder (REC). The fibre-optic microprobe part of the sensor is connected to a photomultiplier detector unit (PM), which is connected to a PC-interfaced controlling unit (PAM control). The photomultiplier detector unit and the LED ring, which served as an actinic light source, were also connected to the controlling unit...129

Appendix A, Figure 2. Schematic illustration of the measuring sequence for obtaining a combined steady-state P vs. actinic irradiance and rETR vs. irradiance curve. Steady-state in levels of oxygen and fluorescence yields were obtained after 10 minutes at each irradiance level (stepped curve). At the end of each 10 minute period a saturating pulse was applied (SP, up arrow) followed by a light dark shift (LD, down arrow). Insert shows the saturating pulse (SP), light-dark shift (LD) and the subsequent increase of irradiance. The sequence was completed within 30 seconds at the end of each 10 minute steady-state acclimation period.....130

Appendix E, Figure 1. Conceptual model based on bleaching sensitivity for branching (*P. damicornis*) and plate-like (*T. reniformis*) corals exposed to 31°C across the GBR. Average sea surface temperature (avr. SST) at the time of collection for each site, degree of bleaching where no bleaching (●), partial bleaching (◐), complete bleaching (○) and zooxanthella type are given.....156

## Table of tables

Table 2.1. Differences in  $F_v/F_m$ , and  $\Phi_{PSII}$  and NPQ of sun- and shade-adapted coenosarc and polyp tissues of *P. damicornis* determined at 358 and 328  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for sun- and shade-adapted coenosarc tissues, respectively, and at 406 and 290  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for sun- and shade-adapted polyp tissues, respectively ( $n = 5 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) between sun- and shade-adapted regions of coenosarc and polyp tissue are denoted with superscript letters. Significant p-value is bolded..... 18

Table 2.2. Quantitative parameters derived from fitted rETR and gross photosynthesis rate curves as a function of PAR. Variable chlorophyll *a* fluorescence:  $rETR_{\text{max}}$  (a.u),  $\alpha_f$ ,  $E_{kf}$  ( $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ); gross photosynthesis rate:  $P_{g\text{max}}$  ( $\text{nmol cm}^{-3} \text{s}^{-1}$ ),  $\alpha_{Pg}$ ,  $E_{kPg}$  ( $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). Values for sun- and shade-adapted coenosarc and polyp tissues are given ( $n = 5 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) between sun- and shade-adapted regions of coenosarc and polyp tissue are denoted with superscript letters. Significant p-values are bolded..... 19

Table 3.1. Observed SSCP genotype frequencies ( $n$ ) of clade A, C1, and C2 ( $f_A$ ,  $f_{C1}$  and  $f_{C2}$ , respectively) are shown as well as their relative proportion (%) to total samples of either sun- or shade-adapted polyps. Bolded values denote the number (or proportion) of polyps in which a given type was dominant. .... 35

Table 4.1a-b. Observed frequencies of *Symbiodinium* type(s) within (sun- and shade-adapted surfaces [abbreviated as Sun and Shade, respectively]) and among colonies of a) *P. damicornis* and b) *A. valida* from Heron Island. Sample numbers ( $n$ ), observed SSCP genotype frequencies ( $f$ ) and proportion (%) of clade A ( $f_A$ ) and C ( $f_{C1}$ ,  $f_{C2}$ ). Where multiple *Symbiodinium* types were present the dominant frequency is bolded... 50

Table 6.1. Collection sites of *P. damicornis* and *T. reniformis* (see also Fig. 1). Observed frequencies of zooxanthella type(s) present in both coral species are given with dominant clade bolded. Shelf position is given. Sample numbers ( $n$ ) and observed SSCP genotype frequencies of clade A, C1, and D ( $f_A$ ,  $f_{C1}$  and  $f_D$ , respectively)..... 84

Table 6.2. Zooxanthella density (numbers of cells  $\times 10^6 \text{ cm}^{-2}$  surface area) of controls ( $26^\circ\text{C}$ ) prior to experimental heating ( $T = 0:1$  replicate in each of 3 bins of 4 experimental temperature tanks,  $n = 12 \pm \text{SE}$ ), after heating ( $T = 14:1$  replicate in each of 3 bins,  $n = 3 \pm \text{SE}$ ) and after recovery ( $T = 28:1$  replicate in each of 3 bins,  $n = 3 \pm \text{SE}$ ) in northern (Lizard Island), central (Big Broadhurst Reef) and southern GBR (*P. damicornis*, Wistari Reef; *T. reniformis*, Heralds Prong Reef). Significant differences ( $p < 0.05$ ) are denoted with asterisks (\*). Significant p-values are bolded. .... 92

Table 6.3. Absolute photochemical efficiency ( $F_v/F_m$ ) ( $n = 27 \pm \text{SE}$ ) prior to experimental heating of explants in northern (Lizard Island), central (Big Broadhurst Reef) and southern GBR (*P. damicornis*, Wistari Reef; *T. reniformis*, Heralds Prong Reef). Significant differences ( $p < 0.01$ ) are denoted with superscript values. Significant p-values is bolded..... 95

Appendix B, Table 1a-b. Quantitative parameters derived from fitted steady-state relative electron transport rates (rETR) and gross photosynthesis rate curves of



individual polyps of *Acropora valida*. Chlorophyll *a* fluorescence— $rETR_{max}$  (a.u),  $\alpha_f$ ,  $E_{kf}$  ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ); gross photosynthesis rate— $P_{gmax}$  ( $\text{nmol cm}^{-3} \text{ s}^{-1}$ ),  $\alpha_{Pg}$ ,  $E_{kPg}$  ( $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ). a) sun- and shade-adapted comparison, b) clade A + C and clade C comparison. Sample size (n) averages and standard errors are given. Tukey's HSD comparison of chlorophyll *a* fluorescence parameters with those of RLC (Appendix B, Table 2a-b) are given as superscript letters.....131

Appendix B, Table 2a-b. Quantitative parameters ( $rETR_{max}$  [a.u],  $\alpha$ ,  $E_k$  [ $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ]) of RLCs at different irradiances derived from fitted relative electron transport rates ( $rETR$ ). Averages and standard errors are given as well as p-value for Mann-Whitney U test (MWH). a) sun- and shade-adapted comparison, b) clade A + C and clade C comparison. P-value (ANOVA) and Tukey's HSD comparisons of RLCs at different irradiances are given as superscript letters and are comparable to those given for SSLCs in Table 1a-b, Appendix B. Significant p-values are bolded.....132

Appendix C, Table 1a-b. Symbiont cell density. Univariate analyses of a) *P. damicornis* and b) *A. valida* of symbiont cell density. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....134

Appendix C, Table 2a-b.  $Dt/(Dn + Dt)$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of  $Dt/(Dn + Dt)$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold. Dn, diadinoxanthin; Dt, diatoxanthin. Dn, diadinoxanthin; Dt, diatoxanthin.....135

Appendix C, Table 3a-b. Non-photochemical quenching,  $NPQ_{RLC}$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of non-photochemical quenching,  $NPQ_{RLC}$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....136

Appendix C, Table 4a-b. Effective quantum yield,  $\Phi_{PSII}$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of effective quantum yield,  $\Phi_{PSII}$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold. ....137

Appendix C, Table 5a-b. Maximum quantum yield,  $F_v/F_m$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of maximum quantum yield,  $F_v/F_m$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold. ....138

Appendix C, Table 6a-b. Maximum quantum yield,  $F_v/F_m$  for sun-adapted fragments Summary of repeated measures ANOVAs comparing responses of maximum quantum yield,  $F_v/F_m$ , to differences among seasons and between measurements at dusk and dawn of a) *P. damicornis* and b) *A. valida*. Year (2004, 2005, 2006; within-subject

factor), season (summer, winter; between-subject factor) and time (dusk, dawn; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....139

Appendix C, Table 7a-b. Maximum quantum yield,  $F_v/F_m$  for shade-adapted fragments Summary of repeated measures ANOVAs comparing responses of maximum quantum yield,  $F_v/F_m$ , to differences among seasons and between measurements at dusk and dawn of a) *P. damicornis* and b) *A. valida*. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and time (dusk, dawn; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....140

Appendix C, Table 8a-b. Quantitative parameters derived from fitted relative electron transport rate,  $rETR$ , curves as a function of PAR. Variable chlorophyll *a* fluorescence:  $rETR_{max}$  (a.u),  $\alpha$ ,  $E_k$  ( $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ); Values for sun- (Sun) and shade-adapted (Shade) a) *P. damicornis* and b) *A. valida* seasonally and over three consecutive years are given. Averages  $\pm$  SE are shown ( $n = 10$ ).....141

Appendix C, Table 9a-b. Maximum relative electron transport rate,  $rETR_{max}$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of maximum relative electron transport rate,  $rETR_{max}$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....142

Appendix C, Table 10a-b. Light utilisation coefficient,  $\alpha$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of the light utilisation coefficient,  $\alpha$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....143

Appendix C, Table 11a-b. Minimum saturation irradiance,  $E_k$ . Univariate analyses of a) *P. damicornis* and b) *A. valida* of minimum saturation irradiance,  $E_k$ . Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....144

Appendix C, Table 12a-b. Fluorescence rise step O. Univariate analyses of a) *P. damicornis* and b) *A. valida* of the fluorescence rise step O. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....145

Appendix C, Table 13a-b. Fluorescence rise step J. Univariate analyses of a) *P. damicornis* and b) *A. valida* of the fluorescence rise step J. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....146

Appendix C, Table 14a-b. Fluorescence rise step I. Univariate analyses of a) *P. damicornis* and b) *A. valida* of the fluorescence rise step I. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position



(sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....147

Appendix C, Table 15a-b. Fluorescence rise step P. Univariate analyses of a) *P. damicornis* and b) *A. valida* of the fluorescence rise step P. Year (2004, 2005, 2006; within-subject factor), season (summer, winter; between-subject factor) and position (sun-adapted, shade-adapted; between-subject factor). Significant differences ( $p < 0.05$ ) are in bold.....148

Appendix D, Table 1a-b. One-way univariate analyses of a) *P. damicornis* and b) *T. reniformis* testing the hypothesis that changes in descriptive parameters,  $P_{\max}$ ,  $\alpha$  and  $E_k$  of fitted PI-curves between winter and summer sampling are independent of location ( $n = 4 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) between sampling time 1 (winter) and 2 (summer) are in bold. LI =Lizard Island, DR =Davies Reef (for *P. damicornis* in summer), BR =Broadhurst Reef (for *P. damicornis* in winter). .....149

Appendix D, Table 2a-b. One-way univariate analyses of a) *Pocillopora damicornis* and b) *Turbinaria reniformis* testing the hypothesis that changes in  $P_{g\max}$  ( $\text{nmol cm}^{-3} \text{ s}^{-1}$ )  $P_{n\max}$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $R_D$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $P_{n\max}:R$  and  $O_{2\max}$  ( $\mu\text{mol l}^{-1}$ ) between winter and summer sampling are independent of location ( $n = 4 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) between sampling time 1 (winter) and 2 (summer) are in bold. LI =Lizard Island, DR =Davies Reef (for *P. damicornis* in summer), BR =Broadhurst Reef (for *P. damicornis* in winter).....150

Appendix D, Table 3a-b. Two-way univariate analyses of a) *P. damicornis* and b) *T. reniformis* testing the hypothesis that changes in descriptive parameters,  $P_{\max}$ ,  $\alpha$  and  $E_k$  of fitted PI-curves following a short-term increase in temperature are independent of location ( $n = 4 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) are in bold. Temperature, in winter: 24°C, 29°C, 32°C, 29°C+32°C; in summer: 29°C, 32°C, 32°C+34°C. Location: LI =Lizard Island, DR =Davies Reef, BR =Broadhurst Reef. ....152

Appendix D, Table 4a-b. Two-way univariate analyses of a) *P. damicornis* and b) *T. reniformis* testing the hypothesis that changes in  $P_{g\max}$  ( $\text{nmol cm}^{-3} \text{ s}^{-1}$ )  $P_{n\max}$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $R_D$  ( $\text{nmol cm}^{-2} \text{ s}^{-1}$ ),  $P_{n\max}:R$  and  $O_{2\max}$  ( $\mu\text{mol l}^{-1}$ ) following a short-term increase in temperature are independent of location. All maximum measurements were obtained at 900  $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$  ( $n = 4 \pm \text{SE}$ ). Significant differences ( $p < 0.05$ ) are in bold. Temperature, in winter: 24°C, 29°C, 32°C, 29°C+32°C; in summer: 29°C, 32°C, 32°C+34°C. Location: LI =Lizard Island, DR =Davies Reef, BR =Broadhurst Reef.....154

Appendix E, Table 1. Univariate analyses of a) heating and, b) recovery of *P. damicornis* testing the hypothesis that changes in  $F_v/F_m$  following a change in heating temperature are independent of sampling origin. Day (d0, d1, d3, d5, d7, d9, d11, d13, d15, d17, d19, d21, d23, d25, d27; within-subject factor), temperature (26°C, 29°C, 31°C, 33°C; between-subject factor) and location (LI, Lizard Island; BBR, Big Broadhurst Reef; WR, Wistari Reef; between-subject factor).). .....156

Appendix E, Table 2. Univariate analyses of a) heating and b) recovery of *T. reniformis* testing the hypothesis that changes in rate of  $F_v/F_m$  recovery following the heating temperature are independent of sampling origin. Day (d0, d1, d3, d5, d7, d9, d11 13,

d15, d17, d19, d21, d23, d25, d27; within-subject factor), temperature (26°C, 29°C, 31°C, 33°C; between-subject factor) and location (LI, Lizard Island; BBR, Big Broadhurst Reef; HPR, Heralds Prong Reef; between-subject factor).....157

## Abstract

Photosynthetic capacity of scleractinian corals relies predominantly on the productivity of single-celled endosymbiotic dinoflagellates of the genus *Symbiodinium*, known as zooxanthellae, residing intracellularly within coral endoderm tissue. The regulation of photosynthesis of zooxanthellae is in turn dependent on light and temperature. This thesis explores the genetic basis for variation in photosynthesis capacity of zooxanthellae by examining the photo-physiology of genetically characterised *Symbiodinium* communities at a range of spatial and temporal scales. *In situ* and manipulative experiments were conducted to improve our understanding of metabolic responses of zooxanthellae under climate change scenarios.

Fine scale measurements of irradiance and photosynthesis allowed the assessment of photo-physiological changes across individual colonies of *Pocillopora damicornis* and *Acropora valida*. *Pocillopora damicornis* generally contain genetically homogeneous populations of *Symbiodinium*, whilst genetically diverse *Symbiodinium* communities exist within *Acropora valida*. Measurements of light absorption in *P. damicornis* were conducted using a scalar irradiance microprobe and it was found that light absorption was greatest in shade-adapted polyp tissue and smallest in sun-adapted coenosarc tissue. Genetic heterogeneities, found at the scale of individual polyps in *A. valida*, correlated with O<sub>2</sub> concentration at the surface of the colony which was greater in polyps that harboured the two clades (A + C) than in polyps that only harboured clade C. In both corals, measurements using an O<sub>2</sub> microelectrode and a fibre-optic microprobe yielded dissimilar results when used at moderate to high irradiances.

Seasonal changes in photosynthetic capacity suggested that *P. damicornis* is more sensitive to combined effects of relatively higher temperature and irradiance in summer than *A. valida* suggesting that the symbiont community of *A. valida* may not be physiologically compromised possibly due to phylogenetic changes of *Symbiodinium*. Furthermore, thermal tolerances of conspecific corals were examined at narrow and wide spatial scales across the length of the Great Barrier Reef. *Pocillopora damicornis*, which harboured *Symbiodinium* type C1, thus bleached in correlation with latitude, whereas *Turbinaria reniformis* bleached in correlation with the presence and absence of the known thermo-tolerant *Symbiodinium* clade D.

The results, integrating over spatial and temporal scales suggest that the acclimatisation capacity of corals to light and temperature is determined by i) history of

light and temperature exposure and in cases where corals associate with multiple *Symbiodinium* types ii) the distribution of *Symbiodinium*.